Thomas Moore Feed™

## SMARTER FEEDING. BETTER RESULTS.




## NEW & IMPROVED FORMULA

# BIRD GUIDE



### BLUE JAY

Has a pointed crest, black ring around its neck and is among the larger of the songbirds. Most comonly seen in deciduous and mixed forests, mixed woodlands, backyards, and parks.



### NORTHERN CARDINAL

Bright red in color, black face, and pointed crest on its head. The female is browner in color. Most comonly seen in woodlands, gardens, parks, and backyards



### CAROLINA CHICKADEE

Can be identified by its tiny stature and its pitch black throat and cap. Most comonly seen in deciduous forests, suburbs, parks, and backyards.



### HOUSE FINCH

Brownish in color with slight white stripes, the male has a red face, neck, and rump. Most comonly seen in urban centers, suburban areas, backyards, edges, yards, and parks.



### CAROLINA WREN

Brown in color with a white breast and stripe above its eyes across the each side of its head. Most comonly seen in deciduous forests, suburbs, parks, and backyards.



### PAINTED BUNTING

One of the most easily noticed birds in Texas, it has a blue head with a red breast, yellow back and green and blue wings. Most commonly seen in forest edges, hedgerows, and thickets.



### WHITE WING DOVE

Easily distinguished from the Mourning Dove by the white stripe on the side of its wings. It is light brown in color with a chunky breast. Most commonly seen in scrub, woodlands, deserts, urban areas, and cultivated areas



### DOWNY WOODPECKER

With a black and white stiped head and a white spotted back can be distinguished among the woodpeckers by the tiny red spot on the back of its head. Most comonly seen in wooded areas.



### TUFTED TITMOUSE

Small grey bird with a pointy crest, black forehead and white breast. Most commonly seen in deciduous woods, mixed evergreen-deciduous woods, orchards, parks, and suburban areas..



### GOLDFINCH

Vibrant yellow in color, has black on his forehead and black wings with a white rump. Most commonly found in open habitats, wooded areas, lowlands and thickets, and urban areas.



### WHITE BREAST NUTHATCH


Grey in color with a white face and belly and has black on the top of its head and down the back of its neck. Most com monly found in mature woods, decidu ous forests, and mixed woodlands..



### HOUSE SPARROW

Brown head with a white belly and brown striped wings. Most comonly seen in urban centers, suburban areas, backyards, edges, yards, and parks.

# FEED BREAKDOWN

| | JAYS | CARDINALS | CHICKADEE | DOVE | GROSBEAK | GOLD FINCH | FINCH | JUNCO | WOODPECKER | TITMOUSE | NUNTHATCH |
|---|---|---|---|---|---|---|---|---|---|---|---|
| |  |  |  |  |  |  |  |  |  |  |  |
|  | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | | ✓ | ✓ | ✓ |
|  | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | | ✓ | ✓ |
|  | ✓ | ✓ | | ✓ | | ✓ | ✓ | ✓ | | ✓ | |
|  | ✓ | ✓ | ✓ | ✓ | | | | ✓ | | ✓ | |
|  | ✓ | ✓ | ✓ | ✓ | | ✓ | ✓ | | | ✓ | |
|  | ✓ | ✓ | ✓ | | ✓ | ✓ | | ✓ | | ✓ | ✓ |
|  | ✓ | ✓ | ✓ | | | ✓ | ✓ | | | ✓ | ✓ |
|  | ✓ | ✓ | ✓ | | | ✓ | ✓ | | | ✓ | ✓ |